

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/340920973>

# COVID-19 pandemic – A focused review for clinicians

Article in *Clinical Microbiology and Infection* · April 2020

DOI: 10.1016/j.cmi.2020.04.023

CITATION

1

READS

1,061

3 authors, including:



Muge Cevik

University of St Andrews

45 PUBLICATIONS 112 CITATIONS

SEE PROFILE



Antonia Ho

University of Glasgow

27 PUBLICATIONS 318 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



The use of CMV QuantiFERON assay in solid organ transplant patients [View project](#)

All content following this page was uploaded by [Antonia Ho](#) on 27 April 2020.

The user has requested enhancement of the downloaded file.

# Journal Pre-proof

COVID-19 pandemic – A focused review for clinicians

Muge Cevik, Connor Bamford, Antonia Ho

PII: S1198-743X(20)30231-7

DOI: <https://doi.org/10.1016/j.cmi.2020.04.023>


Reference: CMI 2024

To appear in: *Clinical Microbiology and Infection*

Received Date: 8 April 2020

Revised Date: 19 April 2020

Accepted Date: 21 April 2020



Please cite this article as: Cevik M, Bamford C, Ho A, COVID-19 pandemic – A focused review for clinicians, *Clinical Microbiology and Infection*, <https://doi.org/10.1016/j.cmi.2020.04.023>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 European Society of Clinical Microbiology and Infectious Diseases. Published by Elsevier Ltd. All rights reserved.

**COVID-19 Pandemic – a focused review for clinicians**

**Author:** Muge Cevik<sup>1</sup>, Connor Bamford<sup>2</sup>, Antonia Ho<sup>3</sup>

**Affiliations:**

1. Division of Infection and Global Health Research, School of Medicine, University of St Andrews, UK

2. Wellcome-Wolfson Institute for Experimental Medicine, School of Medicine, Dentistry and Biomedical Sciences, Queen's University Belfast, Belfast, UK

3. MRC-University of Glasgow Centre for Virus Research, University of Glasgow, Glasgow, UK

**Author of correspondence:**

Name: Dr Muge Cevik

Address: Division of Infection and Global Health Research, School of Medicine, University of St Andrews, Fife, KY16 9TF

Telephone number: 07531993677

Email address: mc349@st-andrews.ac.uk

**Key words:** COVID-19, coronavirus, SARS-CoV-2, novel coronavirus

**Abstract****Background**

The COVID-19 pandemic caused by SARS-CoV-2 remains a significant issue for global health, economics and society. A wealth of data has been generated since its emergence in December 2019 and it is vital for clinicians to keep up with this data from across the world at a time of uncertainty and constantly evolving guidelines and clinical practice.

**Objectives**

Here we provide an update for clinicians on the recent developments about virology, diagnostics, clinical presentation, viral shedding, and treatment options for COVID-19 based on current literature.

**Sources**

We considered published peer-reviewed papers and non-peer-reviewed pre-print manuscripts on COVID19 and related aspects with an emphasis on clinical management aspects.

**Content**

We describe the virological characteristics of SARS-CoV-2 and clinical course of COVID-19 with an emphasis on diagnostic challenges, duration of viral shedding, severity markers and current treatment options.

**Implications**

The key challenge in managing COVID-19 remains the patient density. However, accurate diagnoses as well as early identification and management of high-risk severe cases are important for many clinicians. For improved management of cases, there is a need to understand test probability of serology, qRT-PCR and radiological testing, and the efficacy of available treatment options that could be used in severe cases with a high risk of mortality.

## Introduction

The first cases of atypical pneumonia of unidentified aetiology were reported on December 30, 2019, from Wuhan, China. By January 7, 2020, a novel betacoronavirus, severe acute respiratory syndrome coronavirus (SARS-CoV-2) was identified, while the disease has been named COVID-19. COVID-19 has now been declared a pandemic, affected nearly every country, with over 2.3 million confirmed cases and >160,000 deaths. The initial clinical case series from China largely comprised of hospitalised patients with severe pneumonia. Further data suggested that approximately 80% patients have mild disease, 20% require hospital admission, and approximately 5% require intensive care admission [1]. Mortality rates are higher among people over 60 years and with coexisting conditions; hypertension, diabetes and cardiovascular disease being the most common. Here we provide an update for clinicians on the recent developments about virology, diagnostics, clinical presentation, and treatment options for COVID-19 based on current literature.

## Virology

Metagenomic sequencing and targeted real-time polymerase chain reaction (qRT-PCR) assays identified a novel human CoV (SARS-CoV-2) in bronchoalveolar lavage fluid taken from the initial patient cluster in Wuhan [2]. Infectious SARS-CoV-2 has been cultured on monkey Vero, human Huh7 and primary human airway epithelial cells [3], where it is cytopathic. Furthermore, serum antibodies (IgM and IgG) from cases neutralized SARS-CoV-2 in cell culture and detected virus-infected cells by indirect immunofluorescence [3].

Phylogenetic analysis reveals that SARS-CoV-2 is closely related to SARS-CoV (~80% similar) in the *Sarbecovirus* sub-family (genus *Betacoronavirus*) [2]. While an intermediate host has yet to be determined, it shares strong genetic similarity (>95%) to known bat coronaviruses from China, suggesting a likely bat origin. Relatively similar coronaviruses have been found in pangolins whose receptor-binding domain (RBD) of Spike (S) glycoprotein is more like to SARS-2-CoV-2 than known bat viruses [4].

SARS-CoV-2 shares most of its gene content with SARS-CoV, including the S glycoprotein, the RNA-dependent RNA polymerase (Nsp12) and two proteases papain-like protease (PLpro) and 3C-like protease (3CLpro) [3]. There is also substantial antigenic cross-reactivity between SARS-CoV-2 and SARS-CoV [3, 5]. A recent study confirmed that the angiotensin-converting enzyme 2 (ACE2), expressed in the human respiratory tract epithelium, is the entry receptor for SARS-CoV-2 similar to SARS-CoV and has been shown to cause pneumonia in lab mice only expressing human ACE2 [6, 7]. This is likely mediated by the RBD of the S glycoprotein [8]. Although there is obvious homology between SARS-CoV and SARS-CoV-2, and cross neutralization has been observed [9], significant biological differences, specifically in the S glycoprotein have been noted [5, 10, 11].

#### **Clinical presentation**

A key difference between COVID-19 and seasonal influenza-associated pneumonia is the potential severity of disease even in young adults without comorbidities [12]. In a study that compared three well-conducted Chinese case series to a reference group of patients with influenza-associated pneumonia from 73 German sentinel hospitals, the severity of pneumonia even in adults aged <60 years without chronic preconditions was significantly greater in COVID-19. For instance, 28% of COVID-19 patients treated on the ICU had no reported comorbidity. The rate of ARDS and mechanical ventilation was markedly higher among COVID-19 patients. The median duration of ventilation was 9 days for non-invasive, and 17 days for invasive ventilation [12].

Across all studies, the most common symptoms at onset of illness were fever, cough, fatigue, and myalgia. However, available data suggest that only half of patients are febrile at the time of admission [13, 14]. Gastrointestinal symptoms, including anorexia, nausea, vomiting and diarrhoea are also common, reported in nearly 40% patients in some cohorts [15, 16]. Furthermore, up to 10% patients present with gastrointestinal symptoms without respiratory symptoms or fever [17]. COVID-19 has been associated with a hypercoagulable state with increased risk of venous thromboembolism[18]. Neurological manifestations, including headache, dizziness, altered consciousness, ischaemic and

haemorrhagic strokes, as well as muscle injury, have also been reported [19]. A third of patients reported taste or olfactory disorders in a small Italian cohort, including anosmia [20]. Other extrapulmonary manifestations include skin and ocular manifestations. An Italian study reported cutaneous manifestations in 20% patients [21]. Lastly, ocular manifestations consistent with conjunctivitis was reported in 32% COVID patients in a Chinese case series [22].

The estimated mean incubation period is reported as 3-6 days (range 1.3-11.3) [12]. The duration from symptom onset to dyspnoea was 5-6 days [13, 17] On average, disease progresses further requiring hospitalisation at 7-8 days from symptom onset. Patients may initially appear relatively stable, but they often rapidly deteriorate with severe hypoxia [13, 17]. The key feature seen in these cases is acute respiratory distress syndrome (ARDS) [13, 17]. The interval from symptom onset to the development of ARDS is approximately 8-12 days [13]. In addition, the incidence of cardiovascular manifestations such as myocardial injury seems to be high, likely due to the systemic inflammatory response and immune system disorders during disease progression[23].

Illness severity and development of ARDS are associated with older age and underlying medical conditions [17]. Additionally, neutrophilia, raised lactate dehydrogenase and D-dimer, lymphocyte counts, CD3 and CD4 T-cell counts, AST, prealbumin, creatinine, glucose, low-density lipoprotein, serum ferritin, and prothrombin time were also associated with higher risk of severe disease and ARDS [17]. In a cohort of 191 patients with a definitive clinical outcome (137 discharged and 54 died), mortality was independently associated with older age, higher qSOFA score, d-dimer >1 µg/mL on admission, and the majority had severe disease and experienced complications, such as ARDS, acute kidney injury, and sepsis [13]. Factors most associated with critical illness were admission oxygen saturation <88%, first d-dimer>2500, first ferritin >2500, and first CRP >200 [24]. Furthermore, patients with cardiovascular disease were shown to be more likely to develop severe symptoms[23] in keeping with picture seen in MERS-CoV and SARS.

In comparison, most children appear to have mild disease. Among 1391 asymptomatic and symptomatic children (median age: 6.7 years) with known COVID19 contact in Wuhan Children's Hospital [25], 171 (12.3%) were SARS-CoV2-positive; 27 (15.8%) had no symptoms or radiologic features of pneumonia, 33 (19.3%) had upper respiratory symptoms, and 64.9% had pneumonia. Three patients (with coexisting conditions) required intensive care and 1 death.

In terms of co-infections, a pre-print examining >8000 samples of COVID-19 contacts tested for SARS-CoV2 in China reported viral co-infections in 5.8% of COVID-19 positive individuals (including seasonal coronaviruses, influenza A virus and rhinoviruses [26]. Another study of 1206 patients identified viral co-infection in 24 of 116 (21%) SARS-CoV2-positive patients; rhino/enterovirus, respiratory syncytial virus, and seasonal CoVs were most common [27]. Bacterial and fungal co-infections with SARS-CoV-2 have been documented especially in the ICU setting, including *Acinetobacter baumannii* and *Klebsiella pneumoniae* [28]. Among 191 patients, non-survivors were more likely to have sepsis based on qSOFA score and secondary infection, although detailed bacteriology results were not reported [13]. Secondary infection and positive association between steroid administration and secondary infection should be explored further.

### **Molecular and Serological Diagnosis**

The first genome sequence for SARS-CoV-2 was released on virological.org on 10 January (GenBank accession number MN908947). This allowed the rapid development of several sensitive and specific qRT-PCR assays [29]. Many laboratories worldwide are now able to test for SARS-CoV-2. Assays have been described that detect <10 copies of SARS-CoV-2 per reaction and will not cross-react with SARS-CoV or other human coronaviruses [29]. However, sensitivity and specificity of these tests remain unknown and there is no clear consensus on which is preferred.

Viral RNA loads by qRT-PCR were substantially higher in sputum compared to throat swabs [3, 30, 31], suggesting that the type of sample may also influence the outcome of the test. Therefore, currently submission of both lower and upper respiratory tracts samples is advised.

166

167 Precise molecular detection is hampered by the variability in viral loads in the upper respiratory tract,  
168 especially at later stages of infection. In a study from China, among 241 COVID-19 patients with at  
169 least one positive SARS-CoV-2 qRT-PCR test result, in the first test, 384 (63.0%) were negative [32].  
170 In addition, several tests at different points were variable from the same patients during the course of  
171 diagnosis and treatment.[32]. Therefore, a single positive test should be confirmed by a second qRT-  
172 PCR assay targeting a different SARS-CoV-2 gene. Although, similar studies in Taiwan and Hong  
173 Kong reported less false-negatives[33]. Secondly, a single negative SARS-CoV-2 test (especially if  
174 from upper respiratory tract specimen) or a positive test result for another respiratory pathogen result  
175 should not be used to exclude COVID-19 infection. These findings indicate that qRT-PCR has low  
176 probability of ruling out an infection and in clinically high suspicious cases repeat sampling and also  
177 CT images may need to be used to guide the diagnosis.

178

179 Antibody-based methods to detect seroconversion in serum or plasma based upon enzyme-linked  
180 immunosorbent assays (ELISA), indirect-immunofluorescence or virus neutralisation have been  
181 reported [34-36]. Around 40-50% patients develop an antibody response to SARS-CoV-2 infection  
182 after 7 days, and the majority by 14 days [35, 37]. S1 has been shown to be more specific than S as an  
183 antigen for SARS-CoV-2 in serological diagnosis [36]. The commercial S1 IgG and IgA assays have  
184 lower specificity but IgA showing higher sensitivity [36]. Recently, an ELISA assay based on  
185 detection of recombinant S protein by serum antibodies demonstrated robust and scalable  
186 determination of seroconversion that will facilitate screening of potential exposed individuals for  
187 evidence of past infection [38]. Since seroconversion occurs relatively late in infection, rapid antibody  
188 tests have a limited role in the diagnosis of acute infection; qRT-PCR remains the 'gold standard'.

189

190 There is an ongoing work to understand protective antibody level and immunological marker. Among  
191 175 recovered laboratory-confirmed COVID-19 patients, neutralizing antibodies (NAb) peaked at 10  
192 to 15 days after disease onset. However, approximately 30% failed to develop good level of NAb  
193 titres (ID50: < 500)[39]. In addition, patients who did not generate NABs at the time of discharge did

not develop NABs thereafter. These results highlight that some patients with SARS-CoV-2 would recover without developing high titers of virus-specific NABs. These findings have some implications for vaccine development and also for convalescent plasma treatment as the donor plasma should be titrated before use in passive therapy. There is less information available on the T cell response during SARS-CoV-2 infection and how it correlates with the NAb titres.

#### **Duration of viral shedding and isolation period**

SARS-CoV-2 RNA has been identified by qRT-PCR in respiratory tract samples 1-2 days prior to symptom onset and can persist for 7-12 days in moderate cases, and up to 2 weeks in severe cases [35, 40]. SARS-CoV-2 has also been detected in whole blood [41], saliva [42], faeces [43], and urine [44] by qRT-PCR (Table 1).. In several case series with serial sampling, viral loads were highest soon after symptom onset [35, 45]. Patients with severe COVID-19 had significantly higher viral load and longer period of viral shedding than mild cases [46]. Prolonged viral RNA shedding has been reported from throat swabs up to 37 days among adult patients [13], and in faeces, for over one month after illness onset in children[40, 47]. However, detection of viral RNA by qRT-PCR does not necessarily equate with infectious virus. No live virus was cultured from 9 mild COVID-19 cases beyond day 8 after symptom onset in throat swabs or sputum despite ongoing high viral load [35]. Persistently high levels of RNA were also identified in the stool of the mild cases, but no live virus was cultured [35]. These findings suggest that patients may continue to shed RNA in various samples for a long period, but this does not equate to infectiousness potential (Table 1). This supports the current guidance of 7-14 days self-isolation from symptom onset. Certain hospitals following a protocol to confirm viral clearance prior to transfer out of dedicated COVID-19 wards, however, this may not be required given the prolonged RNA shedding without the evidence of viable virus. However, whether faecal-oral or faecal-respiratory transmission occurs, and the role of shedding in severe cases in transmission requires further exploration.

#### **Transmission patterns**

A review of modelling studies based on Chinese case numbers report a median basic reproduction number ( $R_0$ ) of 2.79 [48], though  $R_0$  as high as 5.7 have been reported [49]. These estimates are substantially higher than the reproduction number for seasonal influenza ( $\sim 1.3$ ) [50], and indicate that control measures would need to prevent  $>60\%$  transmission to stop the epidemic. Of note,  $R_0$  will vary by setting, and can be substantially reduced by countermeasures, as have been observed in China [51].

It is now clear that a significant proportion of individuals with COVID-19 have very mild or no symptoms. Asymptomatic infection at the time of laboratory testing have been reported [52, 53], though a large proportion go on to develop symptoms. For instance, among 55 asymptomatic carriers with positive qRT-PCR for SARS-CoV-2 in pharyngeal swab samples, 14 went on to develop mild, 39 ordinary, and 2 severe COVID-19 [54]. There have been several reports of SARS-CoV-2 transmission from asymptomatic or presymptomatic persons [55, 56], which poses significant challenges to contact tracing. Nevertheless, the relative contribution of asymptomatic or pre-symptomatic transmission on the overall transmission dynamics of the pandemic remains uncertain. Thus, household studies to study secondary human transmission of SARS-CoV-2 and serosurveys to determine the incidence of asymptomatic and subclinical infections are needed.

A further consideration is superspreading events, whereby a small number of cases are responsible for a disproportionate number of secondary cases. This was a feature of both SARS- and MERS-CoV, responsible for multiple nosocomial outbreaks [57, 58]. Several superspreading events have been reported for COVID-19 [17]. Rapid identification and mitigation of these events will be crucial to controlling this pandemic.

#### **Treatment options in clinical trials**

At present, there are no approved antivirals for SARS-CoV-2. Several antivirals that have shown promise against SARS- or MERS-CoV in vitro and in vivo are currently being evaluated in clinical trials for COVID-19. Lopinavir/ritonavir (LPV/r), a protease inhibitor used as an antiretroviral,

showed inconclusive findings for the treatment of SARS, but demonstrated strong *in vitro* and *in vivo* antiviral activity against MERS-CoV when combined with interferon-beta (IFN $\beta$ ) [59]. The first of a number of clinical trials involving LPV/r was recently published [60]. Among 199 seriously ill laboratory-confirmed COVID-19 patients, no significant difference in clinical improvement, mortality or viral clearance was observed between LPV/r (n=99) and standard care (n=100) arms. However, treatment was instituted late in infection; median time from symptom onset to treatment was 13 days, and >40% of patients had undetectable viral load before or during treatment. The results were complicated by the variable use of other treatments, including interferon, glucocorticoids and antibiotics. Of note, day 28 mortality was lower (not significantly) in those with early treatment (19% vs. 27%) and those who received LPV/r also had lower vasopressor and non-invasive ventilation use. Another promising drug is remdesivir, a novel nucleotide analogue that interferes with nsp12 polymerase [61]. It has shown *in vitro* activity against a wide range of RNA viruses including SARS and MERS-CoV [62, 63], and has also demonstrated superior antiviral activity compared to LPV/r-IFN $\beta$  against MERS-CoV in a mouse model [59]. Against SARS-CoV-2, it has shown promising antiviral activity in Vero E6 cells and Huh7 cells [64]. Remdesivir has been given to a small number of patients with severe COVID-19 through compassionate use, however, given the lack of randomisation and control group interpretation of the findings is difficult [65]. There are ongoing RCTs assessing its efficacy and safety in patients with COVID-19 worldwide, and a study in France evaluating its impact on viral shedding in high and moderate risk contacts in confirmed COVID-19 cases (NCT04259892).

Other candidate antivirals are studied in RCTs, including favipiravir and hydroxychloroquine, which has been shown to inhibit virus cell entry *in vitro* [66]. Hydroxychloroquine (HCQ), an analogue of chloroquine, has demonstrated anti-SARS-CoV-2 activity *in vitro* [67]. Among a small open-label non-randomised study, patients treated with HCQ and HCQ + Azithromycin showed viral load reduction compared to controls. However, there has been significant concerns and ethical issues about the content, the ethical approval of the trial and the peer review process prior to publication raised by several physicians and also the International Society of Antimicrobial Chemotherapy. In a small RCT

of HCQ (n=30), there was no change in viral load or clinical outcome after 7 days [68]. Currently, there are 45 trials evaluating chloroquine or HCQ for the treatment and prophylaxis of COVID-19, including multi-centre RCTs in the UK (RECOVERY, ISRCTN50189673), Europe (DisCoVeRy, NCT04315948) and also globally involving >70 countries (SOLIDARITY, ISRCTN83971151).

Host-targeted therapeutic options are also being explored, such as inhibition of human cytokine interleukin-6 (IL-6), the abundance of which has been associated with worse prognosis [69]. A preprint including 21 patients that received Tocilizumab (an IL-6 receptor inhibitor) reported improvement in symptoms, hypoxaemia and CT changes in the majority of patients[70]. There are ongoing RCTs evaluating tocilizumab and sarilumab, also an IL-6 receptor inhibitor. With insufficient evidence of efficacy for any existing treatments, the IDSA recommends that experimental therapies should only be offered to patients in the context of a clinical trial [71].

There is no licenced vaccine to protect against COVID-19. However, a number of experimental candidates are in development with some already in early clinical trials. Most vaccine candidates focus on immunisation with only the S glycoprotein, which is the major target for neutralization antibodies. Candidate vaccines differ in the mode of S delivery and platforms dependent on recombinant protein, mRNA or viral vectored approaches are being tested. Passive immunisation through transfusion of convalescent sera or plasma containing neutralizing antibodies from recovered donors have been reported in several case series, with clinical improvement reported in recipients [72, 73]. Clinical trials evaluating convalescent plasma as treatment for severe COVID-19 are ongoing.

## Conclusion

A wealth of data has been generated already on COVID-19 since early January 2020. Nevertheless, key questions remain regarding understanding the population at risk and age groups, proportion of individuals that have had asymptomatic infections and their transmission potential, endemicity and seasonality of COVID-19, and whether stringent physical distancing measures will be effective in

countries outside China. The main challenge in managing COVID19 remains the patient density, however, accurate diagnoses as well as early identification and management of high-risk severe cases remains a daily battle for many clinicians. For improved management of cases, there is a need to understand test probability of serology, qRT-PCR and radiological testing, and the efficacy of available treatment options that could be used in severe cases with high risk of mortality.

#### **Authors contributions**

MC, CM, AH drafted the first and subsequent versions of the manuscript, and all authors provided critical feedback and contributed to the manuscript.

#### **Financial support and sponsorship**

None

#### **Conflicts of interest**

MC, CB and AT has nothing to disclose.

**Table 1: Transmission routes**

Journal Pre-proof

| Source      | Mode of transmission                                                              | RNA by PCR<br>(Days since onset of<br>symptoms) | Viable virus<br>(Days since onset of<br>symptoms) |
|-------------|-----------------------------------------------------------------------------------|-------------------------------------------------|---------------------------------------------------|
| Nasopharynx | Droplet                                                                           | Up to 37 days                                   | Up to 7 days (in<br>mild cases)                   |
| Sputum      | Droplet / airborne during aerosolize-<br>producing procedures                     | Up to 37 days                                   | Up to 7 days (in<br>mild cases)                   |
| Stool       | No evidence of faecal-oral<br>transmission                                        | > 30 days                                       | Only 1 report;<br>uncertain                       |
| Blood       | No viable virus to date                                                           | Up to 14 days                                   | No                                                |
| Urine       | No viable virus to date                                                           | No                                              | No                                                |
| Conjunctiva | No viable virus to date<br>Macaques with corneal inoculation<br>develop infection | Yes                                             | No                                                |
| Vertical    | No strong evidence of vertical<br>transmission to date                            | No                                              | N/A                                               |

## References

1. Team TNCPERE. The Epidemiological Characteristics of an Outbreak of 2019 Novel Coronavirus Diseases (COVID-19) — China, 2020. Available at: <http://weekly.chinacdc.cn/fileCCDCW/journal/article/ccdcw/newcreate/COVID-19.pdf>. Accessed February 18.
2. Zhu N, Zhang D, Wang W, et al. A Novel Coronavirus from Patients with Pneumonia in China, 2019. *New England Journal of Medicine* **2020**.
3. Zhou P, Yang X-L, Wang X-G, et al. A pneumonia outbreak associated with a new coronavirus of probable bat origin. *Nature* **2020**.
4. Lam TT-Y, Shum MH-H, Zhu H-C, et al. Identifying SARS-CoV-2 related coronaviruses in Malayan pangolins. *Nature* **2020**.
5. Tian X, Li C, Huang A, et al. Potent binding of 2019 novel coronavirus spike protein by a SARS coronavirus-specific human monoclonal antibody. *bioRxiv* **2020**: 2020.01.28.923011.
6. Bao L, Deng W, Huang B, et al. The Pathogenicity of 2019 Novel Coronavirus in hACE2 Transgenic Mice. *bioRxiv* **2020**: 2020.02.07.939389.

7. Gorbalenya AE, Baker SC, Baric RS, et al. <em>Severe acute respiratory syndrome-related coronavirus</em>: The species and its viruses – a statement of the Coronavirus Study Group. bioRxiv **2020**: 2020.02.07.937862.
8. Letko M, Munster V. Functional assessment of cell entry and receptor usage for lineage B  $\beta$ -coronaviruses, including 2019-nCoV. bioRxiv **2020**: 2020.01.22.915660.
9. Hoffmann M, Kleine-Weber H, Schroeder S, et al. SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked by a Clinically Proven Protease Inhibitor. Cell.
10. Lei C, Fu W, Qian K, et al. Potent neutralization of 2019 novel coronavirus by recombinant ACE2-Ig. bioRxiv **2020**: 2020.02.01.929976.
11. Wrapp D, Wang N, Corbett KS, et al. Cryo-EM Structure of the 2019-nCoV Spike in the Prefusion Conformation. bioRxiv **2020**: 2020.02.11.944462.
12. Tolksdorf K, Buda S, Schuler E, Wieler LH, Haas W. Influenza-associated pneumonia as reference to assess seriousness of coronavirus disease (COVID-19). Euro Surveill **2020**; 25(11).
13. Zhou F, Yu T, Du R, et al. Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: a retrospective cohort study. Lancet **2020**; 395(10229): 1054-62.
14. Kluytmans M, Buiting A, Pas S, et al. SARS-CoV-2 infection in 86 healthcare workers in two Dutch hospitals in March 2020. medRxiv **2020**: 2020.03.23.20041913.
15. Liang W, Feng Z, Rao S, et al. Diarrhoea may be underestimated: a missing link in 2019 novel coronavirus. Gut **2020**: gutjnl-2020-320832.
16. Zhang J-j, Dong X, Cao Y-y, et al. Clinical characteristics of 140 patients infected with SARS-CoV-2 in Wuhan, China. Allergy **2020**; n/a(n/a).
17. Wang D, Hu B, Hu C, et al. Clinical Characteristics of 138 Hospitalized Patients With 2019 Novel Coronavirus-Infected Pneumonia in Wuhan, China. JAMA **2020**.
18. Danzi GB, Loffi M, Galeazzi G, Gherbesi E. Acute pulmonary embolism and COVID-19 pneumonia: a random association? European Heart Journal **2020**.

- 365 19. Mao L, Jin H, Wang M, et al. Neurologic Manifestations of Hospitalized Patients With  
366 Coronavirus Disease 2019 in Wuhan, China. *JAMA Neurology* **2020**.
- 367 20. Giacomelli A, Pezzati L, Conti F, et al. Self-reported olfactory and taste disorders in SARS-  
368 CoV-2 patients: a cross-sectional study. *Clinical Infectious Diseases* **2020**.
- 369 21. Recalcati S. Cutaneous manifestations in COVID-19: a first perspective. *Journal of the*  
370 *European Academy of Dermatology and Venereology* n/a(n/a).
- 371 22. Wu P, Duan F, Luo C, et al. Characteristics of Ocular Findings of Patients With Coronavirus  
372 Disease 2019 (COVID-19) in Hubei Province, China. *JAMA Ophthalmology* **2020**.
- 373 23. Zheng Y-Y, Ma Y-T, Zhang J-Y, Xie X. COVID-19 and the cardiovascular system. *Nature*  
374 *Reviews Cardiology* **2020**; 17(5): 259-60.
- 375 24. Petrilli CM, Jones SA, Yang J, et al. Factors associated with hospitalization and critical  
376 illness among 4,103 patients with COVID-19 disease in New York City. *medRxiv* **2020**:  
377 2020.04.08.20057794.
- 378 25. Lu X, Zhang L, Du H, et al. SARS-CoV-2 Infection in Children. *New England Journal of*  
379 *Medicine* **2020**.
- 380 26. Wang M, Wu Q, Xu W, et al. Clinical diagnosis of 8274 samples with 2019-novel  
381 coronavirus in Wuhan. *medRxiv* **2020**: 2020.02.12.20022327.
- 382 27. Kim D, Quinn J, Pinsky B, Shah NH, Brown I. Rates of Co-infection Between SARS-CoV-2  
383 and Other Respiratory Pathogens. *JAMA* **2020**.
- 384 28. Chen N, Zhou M, Dong X, et al. Epidemiological and clinical characteristics of 99 cases of  
385 2019 novel coronavirus pneumonia in Wuhan, China: a descriptive study. *Lancet* **2020**.
- 386 29. Corman VM, Landt O, Kaiser M, et al. Detection of 2019 novel coronavirus (2019-nCoV) by  
387 real-time RT-PCR. *Euro Surveill* **2020**; 25(3).
- 388 30. Chan JF-W, Yuan S, Kok K-H, et al. A familial cluster of pneumonia associated with the  
389 2019 novel coronavirus indicating person-to-person transmission: a study of a family cluster.  
390 *The Lancet* **2020**; 395(10223): 514-23.

31. Han H, Luo Q, Mo F, Long L, Zheng W. SARS-CoV-2 RNA more readily detected in induced sputum than in throat swabs of convalescent COVID-19 patients. *Lancet Infect Dis* **2020**.
32. Li Y, Yao L, Li J, et al. Stability issues of RT-PCR testing of SARS-CoV-2 for hospitalized patients clinically diagnosed with COVID-19. *Journal of Medical Virology* **2020**; n/a(n/a).
33. Chan JF-W, Yip CC-Y, To KK-W, et al. Improved molecular diagnosis of COVID-19 by the novel, highly sensitive and specific COVID-19-RdRp/Hel real-time reverse transcription-polymerase chain reaction assay validated *in vitro* and with clinical specimens. *Journal of Clinical Microbiology* **2020**: JCM.00310-20.
34. Zhou P, Yang X-L, Wang X-G, et al. Discovery of a novel coronavirus associated with the recent pneumonia outbreak in humans and its potential bat origin. *bioRxiv* **2020**: 2020.01.22.914952.
35. Wolfel R, Corman VM, Guggemos W, et al. Virological assessment of hospitalized patients with COVID-2019. *Nature* **2020**.
36. OKBA NMA, Muller MA, Li W, et al. SARS-CoV-2 specific antibody responses in COVID-19 patients. *medRxiv* **2020**: 2020.03.18.20038059.
37. Zhao J, Yuan Q, Wang H, et al. Antibody responses to SARS-CoV-2 in patients of novel coronavirus disease 2019. *medRxiv* **2020**: 2020.03.02.20030189.
38. Amanat F, Nguyen T, Chromikova V, et al. A serological assay to detect SARS-CoV-2 seroconversion in humans. *medRxiv* **2020**: 2020.03.17.20037713.
39. Wu F, Wang A, Liu M, et al. Neutralizing antibody responses to SARS-CoV-2 in a COVID-19 recovered patient cohort and their implications. *medRxiv* **2020**: 2020.03.30.20047365.
40. Cai J, Xu J, Lin D, et al. A Case Series of children with 2019 novel coronavirus infection: clinical and epidemiological features. *Clinical Infectious Diseases* **2020**.
41. Young BE, Ong SWX, Kalimuddin S, et al. Epidemiologic Features and Clinical Course of Patients Infected With SARS-CoV-2 in Singapore. *JAMA* **2020**.
42. To KK-W, Tsang OT-Y, Yip CC-Y, et al. Consistent Detection of 2019 Novel Coronavirus in Saliva. *Clinical Infectious Diseases* **2020**.

43. Wang W, Xu Y, Gao R, et al. Detection of SARS-CoV-2 in Different Types of Clinical Specimens. *JAMA* **2020**.
44. Peng L, Liu J, Xu W, et al. 2019 Novel Coronavirus can be detected in urine, blood, anal swabs and oropharyngeal swabs samples. *medRxiv* **2020**: 2020.02.21.20026179.
45. Zou L, Ruan F, Huang M, et al. SARS-CoV-2 Viral Load in Upper Respiratory Specimens of Infected Patients. *New England Journal of Medicine* **2020**; 382(12): 1177-9.
46. Liu Y, Yan L-M, Wan L, et al. Viral dynamics in mild and severe cases of COVID-19. *The Lancet Infectious Diseases*.
47. Xu Y, Li X, Zhu B, et al. Characteristics of pediatric SARS-CoV-2 infection and potential evidence for persistent fecal viral shedding. *Nature Medicine* **2020**.
48. Liu Y, Gayle AA, Wilder-Smith A, Rocklöv J. The reproductive number of COVID-19 is higher compared to SARS coronavirus. *Journal of Travel Medicine* **2020**; 27(2).
49. Sanche S, Lin YT, Xu C, Romero-Severson E, Hengartner N, Ke R. High Contagiousness and Rapid Spread of Severe Acute Respiratory Syndrome Coronavirus 2. *Emerg Infect Dis* **2020**; 26(7).
50. Biggerstaff M, Cauchemez S, Reed C, Gambhir M, Finelli L. Estimates of the reproduction number for seasonal, pandemic, and zoonotic influenza: a systematic review of the literature. *BMC Infectious Diseases* **2014**; 14(1): 480.
51. Kucharski AJ, Russell TW, Diamond C, et al. Early dynamics of transmission and control of COVID-19: a mathematical modelling study. *The Lancet Infectious Diseases*.
52. Backer JA, Klinkenberg D, Wallinga J. Incubation period of 2019 novel coronavirus (2019-nCoV) infections among travellers from Wuhan, China, 20-28 January 2020. *Euro Surveill* **2020**; 25(5).
53. Ki M. Epidemiologic characteristics of early cases with 2019 novel coronavirus (2019-nCoV) disease in Korea. *Epidemiol Health* **2020**; 42(0): e2020007-0.
54. Wang Y, Liu Y, Liu L, Wang X, Luo N, Li L. Clinical Outcomes in 55 Patients With Severe Acute Respiratory Syndrome Coronavirus 2 Who Were Asymptomatic at Hospital Admission in Shenzhen, China. *The Journal of Infectious Diseases* **2020**.

55. Kimball A, Hatfield KM, Arons M, et al. Asymptomatic and Presymptomatic SARS-CoV-2 Infections in Residents of a Long-Term Care Skilled Nursing Facility - King County, Washington, March 2020. *MMWR Morb Mortal Wkly Rep* **2020**; 69(13): 377-81.
56. Wei WE, Li Z, Chiew CJ, Yong SE, Toh MP, Lee VJ. Presymptomatic Transmission of SARS-CoV-2 - Singapore, January 23-March 16, 2020. *MMWR Morb Mortal Wkly Rep* **2020**; 69(14): 411-5.
57. Shen Z, Ning F, Zhou W, et al. Superspreading SARS events, Beijing, 2003. *Emerg Infect Dis* **2004**; 10(2): 256-60.
58. Kim KH, Tandir TE, Choi JW, Moon JM, Kim MS. Middle East respiratory syndrome coronavirus (MERS-CoV) outbreak in South Korea, 2015: epidemiology, characteristics and public health implications. *J Hosp Infect* **2017**; 95(2): 207-13.
59. Sheahan TP, Sims AC, Leist SR, et al. Comparative therapeutic efficacy of remdesivir and combination lopinavir, ritonavir, and interferon beta against MERS-CoV. *Nat Commun* **2020**; 11(1): 222.
60. Cao B, Wang Y, Wen D, et al. A Trial of Lopinavir-Ritonavir in Adults Hospitalized with Severe Covid-19. *N Engl J Med* **2020**.
61. Agostini ML, Andres EL, Sims AC, et al. Coronavirus Susceptibility to the Antiviral Remdesivir (GS-5734) Is Mediated by the Viral Polymerase and the Proofreading Exoribonuclease. *mBio* **2018**; 9(2): e00221-18.
62. Warren TK, Jordan R, Lo MK, et al. Therapeutic efficacy of the small molecule GS-5734 against Ebola virus in rhesus monkeys. *Nature* **2016**; 531(7594): 381-5.
63. Sheahan TP, Sims AC, Graham RL, et al. Broad-spectrum antiviral GS-5734 inhibits both epidemic and zoonotic coronaviruses. *Sci Transl Med* **2017**; 9(396).
64. Wang M, Cao R, Zhang L, et al. Remdesivir and chloroquine effectively inhibit the recently emerged novel coronavirus (2019-nCoV) in vitro. *Cell Res* **2020**.
65. Grein J, Ohmagari N, Shin D, et al. Compassionate Use of Remdesivir for Patients with Severe Covid-19. *New England Journal of Medicine* **2020**.

66. Keyaerts E, Vijgen L, Maes P, Neyts J, Van Ranst M. In vitro inhibition of severe acute respiratory syndrome coronavirus by chloroquine. *Biochem Biophys Res Commun* **2004**; 323(1): 264-8.
67. Yao X, Ye F, Zhang M, et al. In Vitro Antiviral Activity and Projection of Optimized Dosing Design of Hydroxychloroquine for the Treatment of Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2). *Clin Infect Dis* **2020**: ciaa237.
68. CHEN Jun LD, LIU Li, LIU Ping, XU Qingnian, XIA Lu, LING Yun, HUANG Dan, SONG Shuli, ZHANG Dandan, QIAN Zhiping, LI Tao, SHEN Yinzong, LU Hongzhou. A pilot study of hydroxychloroquine in treatment of patients with common coronavirus disease-19 (COVID-19). *J Zhejiang Univ (Med Sci)* **2020**; 49(1): 0-.
69. Liu T, Zhang J, Yang Y, et al. The potential role of IL-6 in monitoring severe case of coronavirus disease 2019. *medRxiv* **2020**: 2020.03.01.20029769.
70. Gritti G, Raimondi F, Ripamonti D, et al. Use of siltuximab in patients with COVID-19 pneumonia requiring ventilatory support. *medRxiv* **2020**: 2020.04.01.20048561.
71. Bhimraj Adarsh RLM, Amy Hirsch Shumaker, Valery Lavergne\*\*, Lindsey Baden, Vincent Chi-Chung Cheng, Kathryn M. Edwards, Rajesh Gandhi, William J. Muller, John C. O'Horo, Shmuel Shoham, M. Hassan Murad\*\*, Reem A. Mustafa\*\*, Shahnaz Sultan\*\*, Yngve Falck-Ytter\*\*. Infectious Diseases Society of America Guidelines on the Treatment and Management of Patients with COVID-19. *IDSA* **2020**.
72. Duan K, Liu B, Li C, et al. Effectiveness of convalescent plasma therapy in severe COVID-19 patients. *Proceedings of the National Academy of Sciences* **2020**: 202004168.
73. Shen C, Wang Z, Zhao F, et al. Treatment of 5 Critically Ill Patients With COVID-19 With Convalescent Plasma. *JAMA* **2020**.

**Table 1: Transmission routes**

| <b>Source</b>      | <b>Mode of transmission</b>                                                    | <b>RNA by PCR</b><br>(Days since onset of symptoms) | <b>Viable virus</b><br>(Days since onset of symptoms) |
|--------------------|--------------------------------------------------------------------------------|-----------------------------------------------------|-------------------------------------------------------|
| <b>Nasopharynx</b> | Droplet                                                                        | Up to 37 days                                       | Up to 7 days (in mild cases)                          |
| <b>Sputum</b>      | Droplet / airborne during aerosolize-producing procedures                      | Up to 37 days                                       | Up to 7 days (in mild cases)                          |
| <b>Stool</b>       | No evidence of faecal-oral transmission                                        | > 30 days                                           | Only 1 report; uncertain                              |
| <b>Blood</b>       | No viable virus to date                                                        | Up to 14 days                                       | No                                                    |
| <b>Urine</b>       | No viable virus to date                                                        | No                                                  | No                                                    |
| <b>Conjunctiva</b> | No viable virus to date<br>Macaques with corneal inoculation develop infection | Yes                                                 | No                                                    |
| <b>Vertical</b>    | No strong evidence of vertical transmission to date                            | No                                                  | N/A                                                   |